

## Letter to the Editor

Dear Sir:

Formaldehyde has been shown to cause nasal cancer in rats exposed by inhalation. Neoplasia did not develop at locations distant from sites of initial contact, a finding consistent with the chemical's high reactivity and rapid detoxification. The recently published report of the Consensus Workshop on Formaldehyde, however, drew particular attention to small excesses of malignant brain tumors and leukemia among causes of death in several groups of professionals in the U.S. and U.K. who use formaldehyde to preserve human tissues (1). Despite deficits of these diseases in mortality studies of formaldehyde-exposed industrial workers from the same countries, the report implied that the excesses of brain cancer and leukemia in the professional groups may have been occupationally caused.

When evaluating the significance of the mortality experience of occupational groups, it is important to con-

sider the possible effects of social class. Small disease excesses or deficits may have resulted from lifestyle characteristics of the persons under study as reflected by their social class and not from occupational exposures or "the healthy worker effect." As indicated in Table 1, in the U.S. and U.K. mortality from brain cancer and, to a lesser degree, leukemia, has consistently increased with elevation in social class. An analysis of deaths of white males  $\geq 20$  years of age during 1950-1981 in the state of Washington, U.S., noted proportional mortality ratios (PMRs) of 138 and 128 for brain cancer and leukemia, respectively, among professional and technical workers. These compared to PMRs of 96 and 87 in operatives, and 75 and 66 among laborers (2).

The dichotomy between mortality from brain cancer and leukemia in formaldehyde-exposed professionals (social class I) and workers (operatives and laborers,

Table 1. Mortality ratios of men by social class.

| Population description |       |       |         |                    | Social class |     |          |     |       | Ref. |
|------------------------|-------|-------|---------|--------------------|--------------|-----|----------|-----|-------|------|
| Place                  | Race  | Age   | Years   | Ratio <sup>a</sup> | High I       | II  | III      | IV  | Low V |      |
| Brain cancer           |       |       |         |                    |              |     |          |     |       |      |
| Mass                   | White | ≥ 20  | 1971–73 | SMOR               | 164          | 97  | 114      |     | 62    | (3)  |
| Calif.                 | All   | 20–64 | 1949–51 | SMR                | 130          | 127 | 108      | 77  | 58    | (4)  |
| U.S.                   | All   | 20–64 | 1950    | SMR                | 136          | 121 | 109      | 94  | 81    | (5)  |
| U.K.                   | All   | 15–64 | 1970–72 | SMR                | 108          | 101 | 111, 105 | 100 | 92    | (6)  |
| U.K.                   | All   | 65–74 | 1970–72 | PMR                | 225          | 137 | 109, 99  | 85  | 56    | (6)  |
| U.K.                   | All   | 20–64 | 1949–53 | SMR                | 133          | 96  | 104      | 88  | 99    | (6)  |
| U.K.                   | All   | ≥ 65  | 1949–53 | PMR                | 136          | 112 | 105      | 90  | 71    | (6)  |
| U.K.                   | All   | 35–65 | 1930–32 | SMR                | 167          | 92  | 116      | 97  | 66    | (6)  |
| U.K.                   | All   | 20–65 | 1921–23 | CMFR               | 160          | 160 | 120      | 80  | 60    | (6)  |
| Leukemia               |       |       |         |                    |              |     |          |     |       |      |
| Mass.                  | White | ≥ 20  | 1971–73 | SMOR               | 126          | 97  | 108      |     | 89    | (3)  |
| Calif.                 | All   | 20–64 | 1949–51 | SMR                | 104          | 116 | 101      | 86  | 104   | (4)  |
| U.S.                   | All   | 20–64 | 1950    | SMR                | 117          | 100 | 105      | 89  | 98    | (5)  |
| U.K.                   | All   | 15–64 | 1970–72 | SMR                | 113          | 100 | 107, 101 | 104 | 95    | (6)  |
| U.K.                   | All   | 65–74 | 1970–72 | PMR                | 138          | 124 | 108, 98  | 90  | 77    | (6)  |
| U.K.                   | All   | 20–64 | 1949–53 | SMR                | 123          | 98  | 104      | 93  | 89    | (6)  |
| U.K.                   | All   | ≥ 65  | 1949–53 | PMR                | 202          | 115 | 101      | 78  | 74    | (6)  |
| U.K.                   | All   | 20–65 | 1930–32 | SMR                | 152          | 126 | 97       | 95  | 86    | (6)  |

<sup>a</sup>SMR—standardized mortality ratio; PMR—proportional mortality ratio; SMOR—standardized mortality odds ratio; CMFR—comparative mortality figure ratio.

social classes IV and V) may be ascribed at least in part to factors associated with differences in social class. One such factor might be access to and use of medical diagnostic services and surgery. This could account for a trend in cause-specific mortality with social class even in the absence of a true difference in disease incidence, owing to a greater extent of misdiagnosis or lack of diagnosis among the lower socioeconomic strata. Whatever the reasons may be, social class gradients of disease do exist and must be taken into account by occupational epidemiologists.

RICHARD J. LEVINE  
 Department of Epidemiology  
 Chemical Industry Institute of Toxicology  
 P.O. Box 12137  
 Research Triangle Park, NC 27709

## REFERENCES

1. National Center for Toxicological Research. Report on the Consensus Workshop on Formaldehyde. *Environ. Health Perspect.* 58: 323-381 (1984).
2. Milham, S., Jr. Improving occupational standardized proportionate mortality ratio analysis by social class stratification. *Amer. J. Epidemiol.* 121:472-475(1985).
3. Dubrow, R., and Wegman, D. H. Occupational characteristics of cancer victims in Massachusetts, 1971-1973. U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health, Cincinnati. DHHS (NIOSH) Publication No. 84-109, September 1984.
4. Buell, P., Dunn, J. E., Jr., and Breslow, L. The occupational-social class risks of cancer mortality in men. *J. Chron. Dis.* 12: 600-621 (1960).
5. Guralnick, L. Mortality by occupation level and cause of death among men 20 to 64 years of age: United States, 1950. U.S. Department of Health, Education, and Welfare, Public Health Service, National Vital Statistics Division. *Vital Statistics—Special Reports*, Vol 53, No. 5, September 1963.
6. Registrar General. Decennial supplements for England and Wales: occupational mortality—1921-23, 1930-32, 1949-53, 1970-72. Her Majesty's Stationery Office, London, 1927, 1938, 1958, 1978.